

V. M. Bondarenko^{1✉}, M. V. Styopochkina²**ON PROPERTIES OF POSETS OF MM-TYPE (1, 2, 7)**

We calculate the coefficients of transitivity for all posets of MM-type (1, 2, 7) (i.e. posets, which are minimax equivalent to the poset (1, 2, 7)).

Key words: 1-oversupercritical posets, minimax equivalence, coefficient of transitivity, Hasse diagram, group of automorphisms, MM-type.

Introduction. In 1972 P. Gabriel [14] introduced the notion of representation of a finite quiver $Q = (Q_0, Q_1)$ (Q_0 and Q_1 denote the sets of vertices and arrows, respectively) and an integral quadratic form $q_Q : \mathbb{Z}^n \rightarrow \mathbb{Z}$, $n = |Q_0|$, called by him the quadratic Tits form of the quiver Q :

$$q_Q(z) = q_Q(z_1, \dots, z_n) := \sum_{i \in Q_0} z_i^2 - \sum_{i \rightarrow j} z_i z_j,$$

where $i \rightarrow j$ runs through the set Q_1 . He proved that the quiver Q has finite representation type over a field k if and only if its Tits form is positive (i.e., takes positive value on any nonzero vector).

The above quadratic form is naturally generalized to a finite poset $0 \notin S$:

$$q_S(z) = z_0^2 + \sum_{i \in S} z_i^2 + \sum_{i < j, i, j \in S} z_i z_j - z_0 \sum_{i \in S} z_i.$$

In [13] Yu. A. Drozd showed that a poset S has finite representation type if and only if its Tits form is weakly positive (i.e., takes positive value on any nonzero vector with nonnegative coordinates); representations of posets were introduced by L. A. Nazarova and A. V. Roiter [15].

Note that, in contrast to quivers, the sets of posets with weakly positive and with positive Tits form do not coincide. Posets with positive Tits forms were studied by the authors in many papers (see e.g. [1–10]).

In particular, in [6] it is introduced the notion of P -critical poset (a poset S is said to be P -critical if its Tits form is not positive, but that of any proper subset of S is positive). Such posets are analogs of the extended Dynkin diagrams. They are classified in [6].

Combinatorial properties of P -critical posets were studied by the authors in [11]. Analogous properties were studied in [12] for the posets of MM-type (1, 3, 5).

In this paper we continue these investigations.

1. Coefficients of transitivity. In this section we recall the notion of coefficient of transitivity of a poset [11].

Let S be a finite poset and $S_{<}^2 := \{(x, y) \mid x, y \in S, x < y\}$. If $(x, y) \in S_{<}^2(x, y)$ and there is no z satisfying $x < y < z$, then x and y are called neighboring. We put $n_w = n_w(S) := |S_{<}^2|$ and denote by $n_e = n_e(S)$ the number of pairs of neighboring elements. On the language of the Hasse diagram $H(S)$ (that represents S in the plane), n_e is equal to the number of all its edges and n_w to the number of all its paths, up to parallelity, going bottom-up (two path is called parallel if they start and terminate at the same

✉ vitalij.bond@gmail.com

vertices). The ratio $k_t = k_t(S)$ of the numbers $n_w - n_e$ and n_w is called the coefficient of transitiveness of S . In the case $n_w = 0$ (then $n_e = 0$) $k_t = 0$.

2. The posets of MM-type (1, 2, 7). For a fixed poset P we say that a poset S is of the MM-type P if S is (min, max)-equivalent to P (the notion of (min, max)-equivalence was introduced in [1]; see also [6]). Since some time we have been used the term minimax equivalence.

The poset (1, 2, 7) (the disjoint union of chains of length 1, 2 and 7) is an element with trivial group of automorphisms in the set of 1-oversupercritical posets (see [2]). The posets of MM-type (1, 2, 7) were classified in [2]. They are given (up to isomorphism and anti-isomorphism) by the following table.

B-1	B-2	B-3	B-4	B-5	B-6
B-7	B-8	B-9	B-10	B-11	B-12
B-13	B-14	B-15	B-16	B-17	B-18
B-19	B-20	B-21	B-22	B-23	B-24

3. Main result. In [11] the authors calculate the coefficient of transitiveness for all P -critical posets. Analogous result was obtained in [12] for the posets of MM-type (1, 3, 5). In this paper we do it for the posets of MM-type (1, 2, 7).

We write all the coefficients of transitiveness k_t up to the second decimal place.

Theorem 1. The following holds for posets (B-1)–(B-24) of the MM-type (1, 2, 7):

N	n_e	n_w	k_t	N	n_e	n_w	k_t	N	n_e	n_w	k_t
1	7	22	0.68	9	8	18	0.56	17	9	24	0.63
2	9	31	0.71	10	9	43	0.79	18	9	22	0.59
3	8	29	0.72	11	9	29	0.69	19	9	29	0.69
4	8	19	0.58	12	9	23	0.61	20	9	20	0.55
5	9	36	0.75	13	9	31	0.71	21	8	22	0.64
6	9	26	0.65	14	9	21	0.57	22	9	23	0.61
7	9	38	0.76	15	8	19	0.58	23	9	21	0.57
8	9	24	0.63	16	8	27	0.70	24	8	27	0.70

The proof is carried out by direct calculations.

1. Bondarenko V. M. On (min,max)-equivalence of posets and applications to the Tits forms // Visn. Kyiv Univ., Ser. Fiz. Mat. – 2005. – No. 1. – P. 24–25.
2. Bondarenko V. V., Bondarenko V. M., Stepochkina M. V., Chervyakov I. V. 1-over-supercritical partially ordered sets with trivial group of automorphisms and min-equivalence. I // Nauk. Visn. Uzhgorod. Univ., Ser. Mat. Inf. – 2011. – **22**, No. 2. – P. 17–25 (in Russian).
3. Bondarenko V. M., Polishchuk A. M. On a criterion of positive definiteness for one class of infinite quadratic forms // Nelin. Kolyv. – 2003. – **6**, No. 1. – P. 3–14 (in Russian).
Bondarenko V. M., Polishchuk A. M. On a criterion of positive definiteness for one class of infinite quadratic forms // Nonlinear Oscil. – 2003. – **6**, No. 1. – P. 1–12. – <https://doi.org/10.1023/A:1024886517654>
4. Bondarenko V. M., Polishchuk A. M. On finiteness of critical Tits forms of posets // Proc. Inst. Math. NAS Ukraine. – 2004. – **50**, Part 3. – P. 1061–1063.
5. Bondarenko V. M., Styopochkina M. V. On posets of width two with positive Tits form // Algebra and Discrete Math. – 2005. – **4**, No. 2. – P. 20–35.
6. Bondarenko V. M., Styopochkina M. V. (Min,max)-equivalence of posets and quadratic Tits form // In: Problems of Analysis and Algebra: Zb. Pr. Inst. Mat. NAN Ukr. – 2005. – **2**, No. 3. – P. 18–58 (in Russian).
7. Bondarenko V. M., Styopochkina M. V. On finite posets of inj-finite type and their Tits forms // Algebra Discrete Math. – 2006. – **5**, No. 2. – P. 17–21.
8. Bondarenko V. M., Stepochkina M. V. On serial posets with positive-definite quadratic Tits form // Nelin. Kolyv. – 2006. – **9**, № 3. – P. 320–325 (in Ukrainian).
Bondarenko V. M., Stepochkina M. V. On serial posets with positive-definite quadratic Tits form // Nonlinear Oscil. – 2006. – **9**, No. 3. – P. 312–316. – <https://doi.org/10.1007/s11072-006-0045-y>
9. Bondarenko V. M., Stepochkina M. V. (Min,max)-equivalence of posets and nonnegative Tits forms // Ukr. Mat. Zh. – 2008. – **60**, № 9. – 1157–1167 (in Russian).
Bondarenko V. M., Stepochkina M. V. (min,max)-equivalence of posets and nonnegative Tits forms // Ukr. Math. J. – 2008. – **60**, No. 9. – 1349–1359. – <https://doi.org/10.1007/s11253-009-0147-7>
10. Bondarenko V. M., Stepochkina M. V. Description of posets critical with respect to the nonnegativity of the quadratic Tits form // Ukr. Math. Zh. – 2009. – **61**, № 5. – P. 611–624 (in Russian).
Bondarenko V. M., Stepochkina M. V. Description of posets critical with respect to the nonnegativity of the quadratic Tits form // Ukr. Math. J. – 2009. – **61**, No. 5. – P. 734–746. – <https://doi.org/10.1007/s11253-009-0245-6>

11. Bondarenko V. M., Styopochkina M. V. Coefficients of transitiveness of P -critical posets // In: Analysis and application: Zb. Pr. Inst. Mat. NAN Ukr. – 2017. – **14**, No. 1. – P. 46–51.
12. Bondarenko V. M., Styopochkina M. V. On properties of posets of MM-type (1,3,5) // Nauk. Visn. Uzhgorod. Univ., Ser. Mat. Inf. – 2018. – **32**, No. 1. – P. 50–53.
13. Drozd Yu. A. Coxeter transformations and representations of partially ordered sets // Funkc. Anal. i Priložen. – 1974. – **8**, No. 3. – P. 34–42 (in Russian).
14. Gabriel P. Unzerlegbare Darstellungen I // Manuscripta Math. – 1972. – **6**. – P. 71–103. – <https://doi.org/10.1007/BF01298413>
15. Nazarova L. A., Roiter A. V. Representations of posets // Zap. Nauch. Semin. LOMI. – 1972. – **28**. – P. 5–31 (in Russian).

ПРО ВЛАСТИВОСТІ ЧАСТКОВО ВПОРЯДКОВАНИХ МНОЖИН ММ-ТИПУ (1, 2, 7)

Обчислюємо коефіцієнт транзитивності для всіх частково впорядкованих множин ММ-типу (1, 2, 7) (тобто частково впорядкованих множин, які мінімаксно еквівалентні частково впорядкованій множині (1, 2, 7)).

Ключові слова: 1-надсуперкритична частково впорядкована множина, мінімаксна еквівалентність, коефіцієнт транзитивності, діаграма Хассе, група автоморфізмів, ММ-тип.

О СВОЙСТВАХ ЧАСТИЧНО УПОРЯДОВАННЫХ МНОЖЕСТВ ММ-ТИПА (1, 2, 7)

Вычисляем коэффициент транзитивности для всех частично упорядоченных множеств ММ-типу (1, 2, 7) (то есть частично упорядоченных множеств, которые минимаксно эквивалентны частично упорядоченному множеству (1, 2, 7)).

Ключевые слова: 1-надсуперкритическое частично упорядоченное множество, минимаксная эквивалентность, коэффициент транзитивности, диаграмма Хассе, группа автоморфизмов, ММ-тип.

¹ Institute of Mathematics of NAS of Ukraine, Kyiv

Received

² Zhytomyr Nat. Univ. of Agriculture and Ecology, Zhytomyr

01.09.19